



Editorial

Taxonomy in crisis: Addressing the shortage of taxonomists in a biodiversity hotspot era

Abdulla-Al-Asif^{1,2*} , Sayali Nerurkar³ 



¹ *Research and Development Wing, Genesis Research Consultancy Limited, Navaron, Jadabpur-7432, Jashore, Bangladesh*

² *Department of Animal Science and Fishery, Faculty of Agricultural and Forestry Sciences, University Putra Malaysia, Bintulu Campus, Nyabau Road, 97008 Bintulu, Sarawak, Malaysia*

³ *Ecological Society, Pune, Maharashtra, India*

Article info

Article history

Received: 10 October 2024

Revised: 20 February 2024

Accepted: 26 October 2024

Published: 26 October 2024

Keywords

Species discovery

Ecological conservation

DNA barcoding

Capacity building

Biodiversity loss

Abstract

In the era of biodiversity crises, where the extinction rates are accelerating at an unprecedented pace, taxonomy—the science of classification, description, and identification of organisms—stands as a critical tool for understanding and preserving the natural world. Yet, paradoxically, this field is in crisis. The shortage of trained taxonomists has become a significant bottleneck in biodiversity research and conservation efforts, especially in regions designated as biodiversity hotspots, which are home to the most unique and endangered species on the planet. This editorial explores the implications of this crisis, the challenges facing taxonomy today, and the steps needed to ensure the discipline can continue to play a pivotal role in safeguarding Earth's biodiversity. Taxonomy provides the foundational knowledge necessary for a variety of scientific disciplines and conservation efforts. Accurate species identification is the first step toward understanding ecosystem dynamics, species interactions, and evolutionary relationships. Without a robust taxonomic framework, it becomes impossible to prioritize conservation efforts effectively. Conservation strategies depend heavily on identifying species at risk of extinction, protecting endangered habitats, and managing invasive species—all tasks that rely on precise taxonomic data.

© 2024 Al-Asif and Nerurkar. This is an open access article distributed under the **Creative Commons Attribution 4.0 International License** (www.creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the era of biodiversity crises, where the extinction rates are accelerating at an unprecedented pace, taxonomy—the science of classification, description, and identification of organisms—stands as a critical tool for understanding and preserving the natural world (Cowie *et al.*, 2022; Löbl *et al.*, 2023). Yet, paradoxically, this field is in crisis (Agnarsson and Kuntner, 2007). The shortage of trained taxonomists has become a significant bottleneck in biodiversity research and conservation efforts, especially in regions designated as biodiversity hotspots, which are home to the most unique and endangered species on the planet (Aronne *et al.*, 2023; Engel *et al.*, 2021). This editorial explores the implications of this crisis, the challenges facing taxonomy today, and the steps needed to ensure the discipline can continue to play a pivotal role in safeguarding Earth's biodiversity.

Taxonomy provides the foundational knowledge necessary for a variety of scientific disciplines and conservation efforts (Dubois, 2003; Sandall *et al.*, 2023). Accurate species identification is the first step toward understanding ecosystem dynamics, species interactions, and evolutionary relationships (Kürzel *et al.*, 2022; Ruppert *et al.*, 2019). Without a robust taxonomic framework, it becomes impossible to prioritize conservation efforts effectively. Conservation strategies depend heavily on identifying species at risk of extinction, protecting endangered habitats, and managing invasive species—all tasks that

rely on precise taxonomic data (Ribeiro *et al.*, 2016).

In biodiversity hotspots, where ecosystems are characterized by high species endemism (species found only in specific regions) and extreme vulnerability to anthropogenic pressures, the role of taxonomists is even more crucial (Esperon-Rodriguez *et al.*, 2024; Kier *et al.*, 2009). These regions, such as the Amazon rainforest, the Western Ghats of India, and the Coral Triangle, are home to a large proportion of Earth's biodiversity (Al-Asif *et al.*, 2022; Gardner *et al.*, 2010). Understanding the diversity of life in these areas requires comprehensive taxonomic assessments to identify species that may be new to science, catalog those at risk, and implement conservation measures (Braby and Williams, 2016; Gallagher *et al.*, 2021). Yet, the shortage of taxonomists in these areas creates a vacuum in knowledge, putting many species at risk of going extinct before they are even discovered (Engel *et al.*, 2021; Löbl *et al.*, 2023; Petrović, 2022).

The term "taxonomic impediment" refers to the shortage of taxonomic expertise and the lack of comprehensive taxonomic information, which hinder the progress of biodiversity research and conservation (Ebach *et al.*, 2011; Engel *et al.*, 2021). This problem is exacerbated by a variety of factors including,

a. Aging workforce: Taxonomy, particularly in its traditional form, is seen by many as a declining field. The majority of practicing taxonomists belong to an older generation, with fewer young scientists entering the field. As these taxonomists retire, the loss of accumulated knowledge and expertise is significant, and the gap left behind is not being filled fast enough (Lücking, 2020).

b. Lack of funding: Taxonomy, often considered a "basic" science, receives comparatively less funding than other fields like genomics, medicine, or environmental science. The lack of financial

*Corresponding authors

Email address: jessoreboyhemel@gmail.com (Abdulla-Al-Asif)

doi: <https://doi.org/10.69517/jars.2024.01.02.0001>

support for taxonomic research and training has dissuaded many young scientists from entering the field, as they are drawn to disciplines with more stable career prospects (Tahseen, 2014).

c. Shift in scientific trends: The rise of molecular techniques and the surge in interest in fields such as genomics and bioinformatics have, in some ways, overshadowed traditional taxonomic methods. While molecular taxonomy (e.g., DNA barcoding) offers powerful tools for species identification, it does not replace the need for experts who understand the morphological, ecological, and behavioral aspects of organisms—critical components of species classification (Tsuballa *et al.*, 2023).

d. Educational barriers: Taxonomy as a discipline is not widely emphasized in modern biology curricula. University programs offering specialized courses in taxonomy have dwindled, leaving fewer opportunities for students to receive formal training in the field. Furthermore, taxonomic research often requires extensive fieldwork, access to museum collections, and long-term study—elements that can be challenging to fit into the fast-paced, results-driven environment of modern academia (Irfanullah, 2002; Wilson, 2004).

The consequences of this taxonomic impediment are far-reaching. Biodiversity data becomes incomplete or inaccurate, slowing conservation efforts and policy-making. Moreover, the lack of taxonomists hampers our ability to respond to emerging global challenges, such as the spread of invasive species and the effects of climate change on species distributions. Without precise taxonomic knowledge, many ecosystems could be irreversibly altered before we even understand what is being lost (Dar *et al.*, 2012; Engel *et al.*, 2021; Löbl *et al.*, 2023).

Biodiversity hotspots are defined by two key criteria: they must contain at least 1,500 species of vascular plants as endemics, and they must have lost at least 70% of their original habitat. These regions are home to nearly 50% of the world's plant species and over 42% of terrestrial vertebrates, despite covering only 2.3% of Earth's surface (Kobayashi *et al.*, 2019; Reid, 1998). The biodiversity within these hotspots is under immense pressure from deforestation, urbanization, agricultural expansion, and climate change. The role of taxonomy in these regions cannot be overstated, yet the shortage of taxonomists is felt most acutely here (Aukema *et al.*, 2017).

For example, in the Amazon rainforest—a biodiversity hotspot with thousands of species yet to be described—the shortage of taxonomists has slowed the process of species identification. This not only hampers conservation efforts but also limits our understanding of how environmental changes, such as deforestation, impact the ecosystem as a whole (Decaëns *et al.*, 2018; Guayasamin *et al.*, 2024). The situation is similar in marine biodiversity hotspots, such as coral reefs in the Coral Triangle, where the loss of taxonomic expertise has made it difficult to assess the true impact of coral bleaching and ocean acidification on marine life (Harvey *et al.*, 2018; Huang *et al.*, 2018). Without taxonomists working in these regions, we risk losing species before they are even discovered. Many endemic species, which are found only in specific regions, could vanish due to habitat loss or climate change, and without taxonomic studies, their extinction might go unnoticed (Lees and Pimm, 2015; Löbl *et al.*, 2023). While the challenges facing taxonomy are significant, there are innovative solutions that can help address the current crisis. Some of these approaches involve leveraging new technologies, while others focus on reinvigorating traditional taxonomic methods and promoting collaboration across disciplines.

1. DNA barcoding and genomic tools: DNA barcoding, a technique that uses a short genetic sequence from a standardized region of the genome to identify species, has revolutionized taxonomy. This method allows for the rapid identification of species, even from minimal or degraded samples, which is particularly useful in biodiversity hotspots where many species are difficult to collect or observe in the wild. Coupling DNA barcoding with next-generation sequencing technologies can help overcome some of the bottlenecks in species identification. However, these tools should complement

rather than replace traditional taxonomic methods (Kress and Erickson, 2008; Odah, 2023).

2. Citizen science and public engagement: Citizen Science initiatives have gained popularity as a way to involve the public in scientific research. In taxonomy, platforms such as iNaturalist and eBird have allowed amateur naturalists to contribute valuable data on species observations. Engaging the public in taxonomic research can help bridge the gap created by the shortage of professional taxonomists. Furthermore, increased public awareness of the importance of taxonomy can drive demand for more educational programs and funding for taxonomic research (Aristeidou *et al.*, 2021).

3. Digital taxonomy and artificial intelligence: The use of artificial intelligence (AI) in species identification is another promising area. AI algorithms, trained on large datasets of species images and descriptions, can assist in the identification of species from photographs or specimens. While still in its infancy, digital taxonomy has the potential to streamline the classification process and make it more accessible to researchers in biodiversity-rich regions with limited access to taxonomic expertise (Bartlett *et al.*, 2022).

4. Capacity building and training: Addressing the shortage of taxonomists will require a concerted effort to build capacity in the field. This includes providing more funding for taxonomic research, creating specialized training programs, and encouraging interdisciplinary collaborations between taxonomists, ecologists, and conservationists. Developing taxonomic expertise in biodiversity hotspots should be a priority, with local universities and research institutions playing a key role in training the next generation of taxonomists (Sandall *et al.*, 2023; Smith and Figueiredo, 2009).

5. Global collaboration and data sharing: The creation of global databases such as the Global Biodiversity Information Facility (GBIF) and the Catalogue of Life has made taxonomic data more accessible to researchers worldwide. These platforms promote data sharing and collaboration, allowing taxonomists to work together across borders. Strengthening international collaborations can help mitigate the effects of the taxonomic crisis, particularly in regions where expertise is lacking (Luo *et al.*, 2021; Sandall *et al.*, 2023).

The current shortage of taxonomists presents a serious challenge to biodiversity research and conservation efforts, particularly in biodiversity hotspots where many species are at risk of extinction. The taxonomic crisis is driven by a combination of factors, including an aging workforce, lack of funding, and shifting scientific priorities. However, through the use of new technologies, public engagement, capacity building, and international collaboration, it is possible to address this crisis and ensure that taxonomy remains a vital tool for understanding and preserving the natural world. As we move further into the Anthropocene—a geological epoch defined by human impact on Earth's ecosystems—the need for taxonomic expertise has never been greater. It is essential that we recognize the importance of taxonomy in biodiversity conservation and take the necessary steps to support and revitalize this critical field. The future of many species, particularly those in biodiversity hotspots, depends on it.

Acknowledgments

The first author gratefully acknowledges the logistical and financial support provided by the Research and Development division of Genesis Research Consultancy Limited.

Ethical approval statement

None to declare.

Data availability

Not applicable.

Informed consent statement

Not applicable.

Conflict of interest

The authors declare no competing interests.

Authors' contribution

Abdulla-Al-Asif and **Sayali Nerurkar** contributed to the conceptualization and writing of this editorial. Both authors has read and approved the final version of the published editorial.

References

- Agnarsson I and Kuntner M, 2007. Taxonomy in a changing world: Seeking solutions for a science in crisis. *Systematic Biology*, 56(3): 531–539. <https://doi.org/10.1080/10635150701424546>
- Al-Asif A, Abu Hena MK, Hamli H, Idris MH, Gerusu GJ, Ismail J, Bhuiyan MKA, Abualreesh MH, Musa N, Wahid MEA and Mishra M, 2022. Status, biodiversity, and ecosystem services of seagrass habitats within the Coral Triangle in the Western Pacific Ocean. *Ocean Science Journal*, 57(2): 147–173. <https://doi.org/10.1007/s12601-022-00068-w>
- Aristeidou M, Herodotou C, Ballard HL, Higgins L, Johnson RF, Miller AE, Young AN and Robinson LD, 2021. How do young community and citizen science volunteers support scientific research on biodiversity? The case of iNaturalist. *Diversity*, 13(7): 318. <https://doi.org/10.3390/d13070318>
- Aronne G, Fantinato E, Strumia S, Santangelo A, Barberis M, Castro S, Cogoni D, Evju M, Galloni M, Glasnović P, Klisz M, Kull T, Lanfranco S, Lazarević M, Petanidou T, Puchałka R, Ranalli R, Stefanaki A, Surina B and Fišer Ž, 2023. Identifying bottlenecks in the life cycle of plants living on cliffs and rocky slopes: Lack of knowledge hinders conservation actions. *Biological Conservation*, 286: 110289. <https://doi.org/10.1016/j.biocon.2023.110289>
- Aukema JE, Pricope NG, Husak GJ and Lopez-Carr D, 2017. Biodiversity areas underthreat: overlap of climate change and population pressures on the world's biodiversity priorities. *PLoS One*, 12: e0170615. <https://doi.org/10.1371/journal.pone.0170615>
- Bartlett P, Eberhardt U, Schütz N and Beker HJ, 2022. Species determination using AI machine-learning algorithms: Hebeloma as a case study. *IMA Fungus*, 13: 13. <https://doi.org/10.1186/s43008-022-00099-x>
- Braby MF and Williams MR, 2016. Biosystematics and conservation biology: critical scientific disciplines for the management of insect biological diversity. *Austral Entomology*, 55: 1–17. <https://doi.org/10.1111/aen.12158>
- Cowie RH, Bouchet P and Fontaine B, 2022. The sixth mass extinction: fact, fiction or speculation? *Biological Reviews*, 97(2): 640–663. <https://doi.org/10.1111/brv.12816>
- Dar GH, Khuroo AA, Reddy CS and Malik AH, 2012. Impediment to taxonomy and its impact on biodiversity science: an Indian perspective. *Proceedings of the National Academy of Sciences, India Section B: Biological Sciences*, 82(2): 235–240. <https://doi.org/10.1007/s40011-012-0031-3>
- Decaëns T, Martins MB, Feijoo A, Oszwald J, Dolédec S, Mathieu J, Arnaud de Sartre X, Bonilla D, Brown GG, Cuellar Criollo YA, Dubs F, Furtado IS, Gond V, Gordillo E, Le Clec'h S, Marichal R, Mitja D, de Souza IM, Praxedes C and Lavelle P, 2018. Biodiversity loss along a gradient of deforestation in Amazonian agricultural landscapes. *Conservation Biology*, 32(6): 1380–1391. <https://doi.org/10.1111/cobi.13206>
- Dubois A, 2003. The relationships between taxonomy and conservation biology in the century of extinctions. *Comptes Rendus. Biologies*, 326(S1): 9–21. [https://doi.org/10.1016/S1631-0691\(03\)00022-2](https://doi.org/10.1016/S1631-0691(03)00022-2)
- Ebach MC, Valdecasas AG and Wheeler QD, 2011. Impediments to taxonomy and users of taxonomy: accessibility and impact evaluation. *Cladistics*, 27(5): 550–557. <https://doi.org/10.1111/j.1096-0031.2011.00348.x>
- Engel MS, Ceriaco LMP, Daniel GM, Dellapé PM, Löbl I, Marinov M, Reis RE, Young MT, Dubois A, Agarwal I, Lehmann AP, Alvarado M, Alvarez N, Andreone F, Araujo-Vieira K, Ascher JS, Baëta D, Baldo D, Bandeira SA and Zacharie CK, 2021. The taxonomic impediment: a shortage of taxonomists, not the lack of technical approaches. *Zoological Journal of the Linnean Society*, 193(2): 381–387. <https://doi.org/10.1093/zoolinnean/zlab072>
- Esperon-Rodriguez M, Correa-Metrio A, Beaumont LJ, Baumgartner JB and Lenoir J, 2024. Evaluating the impact of protected areas in lowering extinction risks in a biodiversity hotspot. *Biological Conservation*, 297: 110728. <https://doi.org/10.1016/j.biocon.2024.110728>
- Gallagher RV, Butt N, Carthey AJR, Tulloch A, Bland L, Clulow S, Newsome T, Dudaniec RY and Adams VM, 2021. A guide to using species trait data in conservation. *One Earth*, 4(7): 927–936. <https://doi.org/10.1016/j.oneear.2021.06.013>
- Gardner TA, Barlow J, Sodhi NS and Peres CA, 2010. A multi-region assessment of tropical forest biodiversity in a human-modified world. *Biological Conservation*, 143(10): 2293–2300. <https://doi.org/10.1016/j.biocon.2010.05.017>
- Guayasamin JM, Ribas CC, Carnaval AC, Carrillo JD, Hoorn C, Lohmann LG, Riff D, Ulloa UC and Albert JS, 2024. Evolution of Amazonian biodiversity: a review. *Acta Amazonica*, 54(spe1): 1–34. <https://doi.org/10.1590/1809-4392202103601>
- Harvey BJ, Nash KL, Blanchard JL and Edwards DP, 2018. Ecosystem-based management of coral reefs under climate change. *Ecology and Evolution*, 8(12): 6354–6368. <https://doi.org/10.1002/ece3.4146>
- Huang D, Goldberg EE, Chou LM and Roy K, 2018. The origin and evolution of coral species richness in a marine biodiversity hotspot*. *Evolution*, 72(2): 288–302. <https://doi.org/10.1111/evo.13402>
- Irfanullah HM, 2002. Present trends in plant taxonomy in Bangladesh and its future. *Bangladesh Journal of Plant Taxonomy*, 10: 99–111. <https://doi.org/10.3329/bjpt.v10i1.661>
- Kier G, Kreft H, Lee TM, Jetz W, Ibsch PL, Nowicki C, Mutke J and Barthlott W, 2009. A global assessment of endemism and species richness across island and mainland regions. *Proceedings of the National Academy of Sciences*, 106(23): 9322–9327. <https://doi.org/10.1073/pnas.0810306106>
- Kobayashi Y, Okada K and Mori AS, 2019. Reconsidering biodiversity hotspots based on the rate of historical land-use change. *Biological Conservation*, 233: 268–275. <https://doi.org/10.1016/j.biocon.2019.02.032>
- Kress WJ and Erickson DL, 2008. DNA barcodes: genes, genomics, and bioinformatics. *Proceedings of the National Academy of Sciences*, 105(8): 2761–2762. <https://doi.org/10.1073/pnas.0800476105>
- Kürzel K, Kaiser S, Lörz AN, Rossel S, Paulus E, Peters J, Schwentner M, Martinez AP, Coleman CO, Svavarsson J and Brix S, 2022. Correct species identification and its implications for conservation using Haploniscidae (Crustacea, Isopoda) in Icelandic waters as a proxy. *Frontiers in Marine Science*, 8: 795196. <https://doi.org/10.3389/fmars.2021.795196>
- Lees AC and Pimm SL, 2015. Species, extinct before we know them? *Current Biology*, 25(5): R177–R180. <https://doi.org/10.1016/j.cub.2014.12.017>
- Löbl I, Klausnitzer B, Hartmann M and Krell FT, 2023. The silent extinction of species and taxonomists—an appeal to science policymakers and legislators. *Diversity*, 15(10): 1053. <https://doi.org/10.3390/d15101053>
- Lücking R, 2020. Three challenges to contemporaneous taxonomy from a lichen-mycological perspective. *Megataxa*, 1: 78–103. <https://doi.org/10.11646/megataxa.1.1.16>

- Luo M, Xu Z, Hirsch T, Aung TS, Xu W, Ji L, Qin H and Ma K, 2021. The use of Global Biodiversity Information Facility (GBIF)-mediated data in publications written in Chinese. *Global Ecology and Conservation*, 25: e01406. <https://doi.org/10.1016/j.gecco.2020.e01406>
- Odah MAA, 2023. Unlocking the genetic code: Exploring the potential of DNA barcoding for biodiversity assessment. *AIMS Molecular Science*, 10(4): 263–294. <https://doi.org/10.3934/molsci.2023016>
- Petrović A 2022. Sizing the knowledge gap in taxonomy: The last dozen years of Aphidiinae research. *Insects*, 13(2): 170. <https://doi.org/10.3390/insects13020170>
- Reid WV, 1998. Biodiversity hotspots. *Trends in Ecology and Evolution*, 13(7): 275–280. [https://doi.org/10.1016/S0169-5347\(98\)01363-9](https://doi.org/10.1016/S0169-5347(98)01363-9)
- Ribeiro J, Colli GR, Caldwell JP and Soares AMVM, 2016. An integrated trait-based framework to predict extinction risk and guide conservation planning in biodiversity hotspots. *Biological Conservation*, 195: 214–223. <https://doi.org/10.1016/j.biocon.2015.12.042>
- Ruppert KM, Kline RJ and Rahman MS, 2019. Past, present, and future perspectives of environmental DNA (eDNA) metabarcoding: A systematic review in methods, monitoring, and applications of global eDNA. *Global Ecology and Conservation*, 17: e00547. <https://doi.org/10.1016/j.gecco.2019.e00547>
- Sandall EL, Maureaud AA, Guralnick R, McGeoch MA, Sica YV, Rogan MS, Booher DB, Edwards R, Franz N, Ingenloff K, Lucas M, Marsh CJ, McGowan J, Pinkert S, Ranipeta A, Uetz P, Wiczorek J and Jetz W, 2023. A globally integrated structure of taxonomy to support biodiversity science and conservation. *Trends in Ecology and Evolution*, 38(12): 1143–1153. <https://doi.org/10.1016/j.tree.2023.08.004>
- Smith GF and Figueiredo E, 2009. Capacity building in taxonomy and systematics. *Taxon*, 58(3): 697–699. <https://doi.org/10.1002/tax.583001>
- Tahseen Q, 2014. Taxonomy-the crucial yet misunderstood and disregarded tool for studying biodiversity. *Journal of Biodiversity and Endangered Species*, 2(3): 128. <https://doi.org/10.4172/2332-2543.1000128>
- Tsaballa A, Kelesidis G, Krigas N, Sarropoulou V, Bagatzounis P and Grigoriadou K, 2023. Taxonomic identification and molecular DNA barcoding of collected wild-growing orchids used traditionally for salep production. *Plants*, 12(17): 3038. <https://doi.org/10.3390/plants12173038>
- Wilson EO, 2004. Taxonomy as a fundamental discipline. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 359(1444): 739–739. <https://doi.org/10.1098/rstb.2003.1440>



Publisher's note

Genesis Publishing Consortium Limited pledges to maintain a neutral stance on jurisdictional claims shown in published maps and institutional affiliations.
