



Original Research

Synergistic antimicrobial potential of garlic (*Allium Sativum*), black cumin (*Nigella Sativa*), and honey against zoonotic bacteria and fungus: An *in vitro* study from Ethiopia

Hafiz Esmael¹ , Abdu Mohamed¹ , Md. Zaminur Rahman² , Md. Mosharraf Hossen³ , Md. Ashiquen Nobil² , Kazi Abdus Sobur^{2*} 

¹ Jimma University College of Agriculture and Veterinary Medicine School of Veterinary Medicine, P.O. Box 307, Jimma, Ethiopia

² Department of Microbiology and Hygiene, Faculty of Veterinary Science, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh

³ Department of Biotechnology, Faculty of Agriculture, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh



Article info

Article history

Received: 04 June 2024

Revised: 28 July 2024

Accepted: 29 September 2024

Published: 01 October 2024

Keywords

Antimicrobial potential
Garlic
Black cumin oil
Honey
Minimum inhibitory concentration (MIC)
Ethiopia

Abstract

The emergence of antibiotic resistance and the evolution of new pathogenic strains have become significant global health challenges in the 21st century, necessitating the exploration of alternative therapeutic options. Herbal remedies, such as garlic, black cumin seed, and honey, have long been used in traditional medicine, but scientific validation of their antimicrobial potential remains crucial. This study aimed to explore the antimicrobial properties of garlic (*Allium sativum*), black cumin seed (*Nigella sativa*), and honey against bacteria and fungi. Various extraction methods, including cold pressing, Soxhlet extraction, hydrodistillation, and cold maceration, were used to obtain bioactive compounds from these natural ingredients. The antimicrobial activity of the extracts was evaluated *in vitro* using the disc diffusion method against *Staphylococcus aureus* (ATCC 25923), *Bacillus cereus* (ATCC 14579), *Escherichia coli* (ATCC 25922), *Salmonella typhi* (ATCC 19430), and *Candida albicans* (ATCC 10237). The results showed inhibition zones ranging from 0 to 35 mm, with black cumin oil demonstrating the highest antibacterial activity, particularly against *E. coli* and *S. aureus*. The minimum inhibitory concentrations (MIC) ranged from 6.25% to 50%. The study concluded that garlic, black cumin seed oil, and honey possess significant antimicrobial properties, either individually or in combination, suggesting their potential as natural alternatives to combat zoonotic bacterial infections and address the growing concern of drug-resistant pathogens. Further research is encouraged to explore their applications in real-world scenarios and develop new, natural antimicrobial therapies.

© 2024 Esmael et al. This is an open access article distributed under the **Creative Commons Attribution 4.0 International License** (www.creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

The rise of antibiotic resistance and the emergence of new pathogenic strains present significant global health challenges, particularly in the 21st century. This crisis underscores the urgent need for alternative therapeutic strategies to combat zoonotic and opportunistic infections. In both developed and developing nations, there is a growing interest in herbal medicine as a safer and more accessible option for disease treatment and prevention (Islam *et al.*, 2024). Approximately 85% of the global population relies on plant-based remedies, with demand increasing as modern pharmaceuticals face growing limitations, including adverse side effects and resistance development (Vaou *et al.*, 2021). In Africa, up to 80% of the population depends on traditional healers, and plant-based remedies play a crucial role in both human and veterinary health (Okaiyeto and Oguntibeju, 2021; Yineger, 2008). Ethiopia, in particular, stands out, with 80% of the human population and over 90% of livestock reliant on plant-derived therapeutics (Abebe and Chane, 2021).

The World Health Organization (WHO) estimates that between

35,000 and 70,000 plant species have been utilized as medicinal agents, representing a significant portion of the estimated 250,000 plant species globally (Saganuwan *et al.*, 2016). However, only 17% of the world's higher plant species have been scientifically investigated for their medicinal potential, leaving a vast, untapped resource for pharmaceutical development (Sen *et al.*, 2015). Ethiopia, one of Africa's biodiversity hotspots, is home to around 6,500 species of flowering plants, including 92 families with known medicinal properties (UNEP, 2018). This biodiversity offers immense potential for the discovery of novel treatments, particularly in the face of multi-drug-resistance cases, which currently cause around 700,000 deaths annually (Baynes, 2016).

Despite significant advancements in modern medicine, the global health community has not introduced any new classes of antibiotics in the past three decades, with no promising candidates in sight (Chang *et al.*, 2015). Consequently, attention has shifted towards traditional herbal remedies, which are increasingly viewed as safer alternatives due to their lower side effects (Balkrishna *et al.*, 2024).

Among the most prominent medicinal plants, garlic (*Allium sativum*) has long been recognized for its antimicrobial properties. Numerous studies have confirmed garlic's efficacy against a range of pathogens, including both gram-positive and gram-negative bacteria, as well as antibiotic-resistant strains (Gabriel *et al.*, 2022; Magryś *et al.*, 2021; Durairaj *et al.*, 2020). In Ethiopia, garlic is widely cultivated in the highland regions, including Adet, Ambo, Debre-work, Sinana,

*Corresponding authors

Email address: soburkazi@yahoo.com (Kazi Abdus Sobur)

doi: <https://doi.org/10.69517/jber.2024.01.02.0004>

and Jimma. Similarly, black cumin seed (*Nigella sativa*), a plant renowned for its therapeutic properties, has been used for centuries across the Middle East, Eastern Europe, Asia, and Africa to treat various ailments (Hannan et al., 2021). In Ethiopia, black cumin seed is extensively grown in regions such as Amara, Oromiya, SNNP, and Gambella (Dessie et al., 2020). Honey, one of the oldest traditional medicines, has been valued for its therapeutic properties and has been documented in ancient medical literature for its antimicrobial and wound-healing activities (Petrossillo, 2008). The WHO reports a rising trend in the use of traditional medicine, particularly in developed countries, driven by the limitations of conventional medicine, the emergence of multi-drug-resistant pathogens, and the adverse effects of chemical drugs (WHO, 2014).

Both Gram-positive and Gram-negative bacteria exhibit drug resistance. The World Health Organization released a list in 2017 that outlined the global priorities for antibiotic research and development. Among these, the most serious and urgent threats that call for increased international attention and quick action were highly resistant Enterobacteriaceae with *Escherichia coli* and *Klebsiella pneumoniae* (carbapenem-resistant and 3rd generation cephalosporin-resistant), as well as *Pseudomonas aeruginosa* (carbapenem-resistant) (Rana et al., 2024; Hossain et al., 2024; Julqarnain et al., 2022).

Given the growing threat of antimicrobial resistance and the limitations of current pharmaceuticals, there is an urgent need to explore and validate the medicinal properties of natural remedies scientifically. The novelty of this study lies in its comprehensive investigation of the synergistic antimicrobial effects of garlic, black cumin seed oil, and honey against specific zoonotic pathogens. Previous research has examined these natural agents individually, but few studies have explored their combined effects, particularly in the context of zoonotic bacteria and fungi. The objectives of this study are to extract and identify the bioactive compounds present in garlic and black cumin seed, compare the yields and antimicrobial activities of different extraction methods, and evaluate the antibacterial and antifungal efficacy of both single and combined extracts of garlic, black cumin seed, and honey against *Staphylococcus aureus*, *Bacillus cereus*, *Escherichia coli*, *Salmonella typhi*, and *Candida albicans* using the disk diffusion method. Ultimately, the study seeks to assess the potential of these natural substances as alternative treatments for bacterial and fungal infections.

By focusing on zoonotic pathogens, this research contributes not only to the field of antimicrobial resistance but also to the broader efforts in promoting the use of medicinal plants for sustainable healthcare solutions. The results of this study could pave the way for the development of novel, plant-based antimicrobial agents.

2. Materials and Methods

2.1 Ethical approval

No ethical approval is required for this study.

2.2 Study area

The study was conducted in Jimma Town which is located 357 km away from Addis Ababa in southwest Ethiopia; and Gumay Woreda which is found in Jimma Zone of Oromia Regional State, and located at 415 km South West of Finfine and 69 km from Jimma town. Study was placed in Jimma University College of Agriculture and Veterinary Medicine School of Veterinary Medicine, Jimma, Ethiopia (Figure 1).

2.3 Processing of ingredients

2.3.1 Cold pressing of *Nigella sativa* oil

Nigella sativa seeds were collected, sieved to remove impurities, and processed using a cold press method at room temperature (25 °C). The oil was separated from the fibers overnight, filtered, and stored in amber glass bottles at room temperature. Ground *Nigella sativa* seeds were subjected to soxhlet extraction using acetone and ethanol as solvents. The extracted oils were concentrated using a rotary evaporator, weighed, and stored in amber glass bottles.

Essential oils were extracted from ground *Nigella sativa* seeds using hydro-distillation. The process involved heating the seeds in distilled water, collecting the steam in a condenser, and separating the essential oil from the water. The oils were stored at 4 °C in amber glass bottles.




Figure 1. Map of the study area.

2.3.2 Cold maceration of garlic

Garlic was crushed, dried for four weeks, and ground into a powder. The powder was macerated in ethanol and acetone, filtered, and the extracts were concentrated and stored in sterile containers at 4 °C.

2.3.3 Honey collection

Honey was collected from beekeepers in Gumay District and filtered through sterile gauze to ensure purity. Different concentrations of honey solutions (100%, 50%, and 25%) were prepared and stored at 4 °C.

2.3.4 Phytochemical screening

The garlic extracts were screened for various phytochemicals, including tannins, saponins, flavonoids, alkaloids, phenols, anthraquinones, and steroids, following established methods.

2.3.5 Microbial strains and culture preparation

Test microorganisms, including *Staphylococcus aureus* (ATCC 25923), *Bacillus cereus* (ATCC 14579), *Escherichia coli* (ATCC 25922), *Salmonella typhi* (ATCC 19430), and *Candida albicans* (ATCC 10237), were obtained from the Biology Department of Jimma University. The organisms were sub-cultured on nutrient agar, standardized, and used for subsequent antimicrobial assays.

2.3.6 Antimicrobial sensitivity (disc diffusion method)

The antimicrobial activity of the extracts was assessed using the disc diffusion method. Sterile discs were impregnated with plant extracts and placed on inoculated agar plates. The zones of inhibition were measured after 24 hours of incubation at 37 °C.

2.3.7 Minimum inhibitory concentration (MIC)

The MIC is the lowest concentration of an extract that inhibits visible bacterial growth (Radojevic et al., 2012). For this study, stock solutions of 50 mg/ml of each extract were prepared, and double dilutions were performed to achieve concentrations of 25 mg/ml, 12.5 mg/ml, and 6.25 mg/ml (CLSI, 2012). Each dilution was mixed with 10 ml of Muller Hinton broth and inoculated with 20 µl of a bacterial suspension adjusted to 0.5 McFarland standards. After incubation at 37 °C for 24 hours, the mixtures were further inoculated on agar plates. The MIC was recorded as the lowest concentration that inhibited bacterial growth. Positive and negative controls were used, and all tests were performed in triplicate.

2.4 Data analysis

Data were recorded in Microsoft Excel and analyzed using SPSS 20. Antimicrobial activity was expressed as mean inhibition zones for the different extracts tested against each bacterial strain. The data

were expressed as mean of antimicrobial activities of the extracts namely: GAE, GEE, BAE, BEE, and HONEY. For each, we were calculated as the mean comparison value of the zone of inhibition obtained against all individual bacteria.

3. Results

3.1 Percentage yield of extract with different method and with different solvent

Black cumin extracted with the soxhlet solvent extraction method yielded 23 g and 19 g extracts from a 63 g sample/300 ml of ethanol and acetone solvent extracted, respectively. Black cumin extracted with cold pressing extraction method yielded 12 g extracts from 100 g sample. Black cumin extracted with hydro distillation extraction method yielded 1 g extracts from 100g sample /1000 ml of water. Garlic extracted with cold maceration extraction method yielded 6g and 3.7 g extracts from 100 g sample/500 ml of ethanol and acetone solvent extracted respectively (Table 1).

Table 1. Results of extracted garlic and black cumin via different method and different solvent.

Method	Sample	Weight (g)	Solvent	Amount (ml)	Result (g)	Type
Soxhlet	black cumin	63	Ethanol	300	23.3	fixed oil
Soxhlet	black cumin	63	Acetone	300	19	fixed oil
Hydrodistillation	black cumin	100	Water	1500	1	pure oil
Cold press	black cumin	100	None	0	12	pure oil
Cold maceration	Garlic	100	Ethanol	500	6	fixed pellet (wet)
Cold maceration	Garlic	100	Acetone	500	3.7	fixed pellet (wet)

3.2 Phytochemical analysis

The phytochemical screening of the samples revealed the presence of several bioactive compounds. Tannins were detected in the acetone extract with a discoloration reaction using bromine water. Saponins produced a frothy reaction in the ethanol extract but were absent in the acetone extract. Flavonoids were present in both ethanol and acetone extracts, indicated by a yellow color change with HCl. Phlobatannins, phenols, glycosides, and steroids were also present in both extracts, showing red precipitation, dirty green, reddish-brown, and green bluish colors respectively. Alkaloids were detected in the ethanol extract with an orange-red color change but were absent in the acetone extract. However, anthraquinones and terpenoids were not detected in either extract, as indicated by the absence of pink coloration and red-violet color, respectively (Table 2).

Table 2. Phytochemical screening.

Phyto chemical	Reagent	Color change	Ethanol	Ac
Tannin	Bromine water	Discoloration	+	+
Saponin	Distilled water	Frothy	+	-
Flavonoid	HCL	Yellow	+	+
Anthraquinone	HCL, Chloroform, NH4	Pink coloration	-	-
Phlobataminin	HCL	Red precipitation	+	+
Alkaloid	H ₂ SO ₄ , HEAT, MAYER'SRGNT	Orange red	+	-
Phenol	Fe ₃ Cl	Dirty green	+	+
Glycoside	Fe ₃ Cl, H ₂ SO ₄	Reddish-brown	+	+
Terpenoid	CHCl ₃ , H ₂ SO ₄	Red violet color	-	-
Steroid	CHCl ₃ , H ₂ SO ₄	Green bluish	+	+

3.3 Antimicrobial sensitivity test

The antimicrobial activity (measured in mm) of 50% concentrations of black cumin oil, garlic, and honey against the tested strains revealed varying levels of inhibition. Black Cumin Ethanol

Extract (BEE) exhibited the most potent activity, especially against *E. coli* and *S. aureus*, with inhibition zones of 35 mm and 34 mm, respectively. Garlic Ethanol Extract (GEE) also showed significant antimicrobial effects, with inhibition zones ranging from 11 mm against *B. cereus* to 20 mm against both *E. coli* and *S. typhi*. Garlic Acetone Extract (GAE) displayed the least antimicrobial activity, with inhibition zones ranging from 0 mm against *S. aureus* and *B. cereus* to 12 mm against *S. typhi*. Honey demonstrated limited activity, with a 7 mm inhibition zone only against *C. albicans*. The mixture of the extracts provided moderate inhibition across all tested strains, while the control group (Gentamycin/Clotrimazole) consistently showed substantial inhibition, with zones ranging from 10 mm against *C. albicans* to 30 mm against *S. aureus* (Figure 2).




Figure 2. Antimicrobial activity (mm) of 50% concentrations of black cumin oil, garlic and honey against tested strains.

The antimicrobial activity (measured in mm) of 25% concentrations of black cumin oil, garlic, and honey against the tested strains demonstrated varying degrees of inhibition. Black Cumin Ethanol Extract (BEE) showed potency against *E. coli* and *S. aureus*, with inhibition zones of 12 mm and 10 mm, respectively. Garlic Ethanol Extract (GEE) also exhibited notable antimicrobial effects, with inhibition zones 17 mm against *C. albicans* and 9 mm against *S. aureus*, *B. cereus* and *E. coli*. In contrast, Garlic Acetone Extract (GAE) and Honey displayed no antimicrobial activity. The mixture of the extracts offered moderate inhibition across all tested strains, while the control group (Gentamycin/Clotrimazole) consistently showed strong inhibition (Figure 3).




Figure 3. Antimicrobial activities (mm) of 25% concentrations of black cumin oil, garlic and honey against tested strains.

The results presented in Table 5 illustrate the synergistic antimicrobial effects of various combinations of garlic ethanol extract (GEE), black cumin acetone extract (BAE), and black cumin oil extract (Bcp). Among the tested pathogens, *Staphylococcus aureus* showed the highest inhibition zone of 17 mm when treated with a mixture of GEE and BCP (GEExBcp), followed by 14 mm with BAExBcp. Similarly, *Bacillus cereus* exhibited the greatest inhibition zone of 16 mm with the GEExBcp combination, while the other combinations demonstrated slightly lower inhibition zones, ranging from 11 to 15 mm. For *Escherichia coli*, the GEE and BAE combinations (GEExBAE

and BEExBCP both resulted in inhibition zones of 12 mm, whereas the GEExBCP combination showed a slightly reduced effect of 11 mm. In contrast, *Salmonella typhi* displayed relatively consistent inhibition zones across all combinations, with values ranging from 9 to 11 mm. The highest inhibition was observed with BEExBCP at 11 mm. *Candida albicans* demonstrated potent inhibition, with the highest zone of 18 mm for the BAExBCP combination, followed by 17 mm for both GEExBAE and GEExBCP. These results indicate that certain combinations, particularly GEExBCP and BAExBCP, exhibit enhanced antimicrobial activity, potentially due to the synergistic effects of the extracts (Figure 4).




Figure 4. Antimicrobial activities of mix potent sample (25gm/ml). The minimum inhibitory concentration (MIC) results, reveal variations in antimicrobial efficacy across different extracts and pathogens. For *Staphylococcus aureus* and *Bacillus cereus*, black cumin oil (Bcp) demonstrated the lowest MIC at 6.25%, while the other extracts, including garlic acetone extract (GAE), garlic ethanol extract (GEE), black cumin ethanol extract (BEE), and black cumin acetone extract (BAE), had MIC values of 12.5%, except for GAE at 25%. Similarly, for *Escherichia coli*, the mixture of extracts exhibited the lowest MIC at 6.25%, with Bcp and other extracts showing MIC values ranging from 12.5% to 25%. In the case of *Salmonella typhi*, the most potent inhibition was observed for GEE and the mixture at 6.25%, while Bcp had a MIC of 13%. Finally, for *Candida albicans*, GEE, BAE, and Bcp demonstrated the most potent activity with a MIC of 6.25%, while other extracts showed higher MICs, including honey (H) at 25%. These findings highlight the superior antimicrobial activity of Bcp and the synergistic mixture of extracts (Figure 5).




Figure 5. Minimum Inhibitory Concentration test against tested strains.

4. Discussion

The extraction efficiency and antimicrobial efficacy of plant-derived compounds are heavily influenced by the extraction method and the solvent used. This study evaluated the yields and antimicrobial activities of garlic and black cumin seed oil extracts, obtained through various extraction techniques, including cold pressing, Soxhlet

extraction, hydrodistillation, and cold maceration, using different solvents.

The yield from cold pressing of black cumin seed (12 g from 100 g) was moderate compared to Soxhlet extraction and hydrodistillation. This result aligns with Benkeblia (2004), who reported that the low heat generated during cold pressing leads to medium yields and moderate antimicrobial activity. The Soxhlet extraction, especially with ethanol, produced the highest yield (23 g from 63 g), likely due to ethanol's greater polarity, which makes it more efficient at extracting bioactive compounds than acetone. This is consistent with findings Benkaci et al. (2012), who reported that black cumin seed oil extraction with ethanol yields higher percentages due to better solubility of the seed's chemical content.

Hydrodistillation, however, yielded only 1% essential oil from black cumin seed, much lower than the other methods (soxhlet extraction and cold pressing). The challenges of oil recovery during hydrodistillation, such as oil escaping with water, contributed to its inefficiency, which contrasts with some studies suggesting that higher temperatures can improve oil yields. This finding is supported by Benkeblia (2004), who noted that low heat during distillation often results in medium or low yields.

Garlic extraction using cold maceration with ethanol also outperformed acetone, producing 6 g of extract compared to 3.7 g with acetone. Ethanol's higher polarity likely contributed to the higher yield, a conclusion supported by Wolde et al. (2018), who reported similar results with more polar solvents. The qualitative phytochemical analysis revealed the presence of several bioactive compounds, such as phenols, saponins, steroids, and flavonoids, in the ethanol extracts, which were either absent or less abundant in the acetone extracts. This is consistent with the findings of Jadon and Dixit (2014), who also noted similar phytochemical profiles in *Allium* species, highlighting their therapeutic potential.

The antimicrobial activity results further demonstrated the effectiveness of ethanol as an extraction solvent. Ethanol-based garlic extracts exhibited the strongest inhibition zones against *E. coli* and *S. typhi* at 50 mg/ml concentrations, with inhibition zones of 22 mm and 20 mm, respectively. In contrast, the acetone extracts showed lower antimicrobial activity, with maximum inhibition zones of 12 mm against *S. typhi*. The superior antimicrobial performance of ethanol extracts can be attributed to their higher concentration of polar bioactive compounds, such as sulfur compounds and phenolics, which are known for their antimicrobial properties (Cushnie and Lamb, 2020).

Black cumin seed oil extracted with ethanol displayed the highest antimicrobial activity, with inhibition zones of 35 mm against *E. coli* and 34 mm against *S. aureus*, outperforming the positive control (gentamycin). Its antifungal activity was also notable, with a 32 mm inhibition zone against *Candida albicans*, compared to 10 mm for clotrimazole, the positive control. These findings are in line with Yimer (2019), who also documented strong antimicrobial properties in black cumin seed oil.

When comparing the overall efficacy, black cumin seed oil extracted with ethanol and acetone was more potent than garlic extracts, and significantly more effective than extracts obtained via hydrodistillation. The combination of garlic extract, black cumin seed oil, and honey exhibited moderate synergistic effects, though the individual extracts, particularly black cumin seed oil, were more effective. This observation is supported by Ncube et al. (2012) and Zhou et al. (2016), who emphasized the importance of synergistic interactions in enhancing the efficacy of herbal remedies.

The minimum inhibitory concentration (MIC) results further demonstrated the potency of these extracts. For instance, black cumin seed oil extracted with ethanol had an MIC of 12.5%, while garlic acetone extracts showed a higher MIC of 25%, indicating lower potency. These findings are consistent with previous studies, such as Hutabarat (2021) and Sadeghian and Ghazvini (2002), which

highlighted the superior efficacy of ethanol extracts in inhibiting bacterial growth.

While this study provides valuable insights into the antimicrobial potential of garlic, black cumin seed oil, and honey, there are several limitations. First, the study was conducted entirely *in vitro*, which may not fully represent how these extracts interact in living organisms. The lack of *in vivo* testing limits the ability to translate these findings into practical therapeutic applications. Second, the exact mechanisms of action of the bioactive compounds were not explored, leaving questions about how these compounds exert their antimicrobial effects. Third, the study focused on a limited number of zoonotic pathogens, and broader testing on additional microbial species would be necessary to confirm the generalizability of the results. Lastly, while the study demonstrated the efficacy of these natural products, potential variations in extract composition due to factors like plant origin and environmental conditions were not considered, which could affect reproducibility.

The study conclusively demonstrates the significant antimicrobial properties of garlic, black cumin seed oil, and honey, individually and in combination, against zoonotic pathogens. The findings underscore the potential of these natural products as alternative treatments to combat bacterial and fungal infections, particularly in light of rising antimicrobial resistance. Despite the promising *in vitro* results, further *in vivo* studies are required to validate these findings in practical, real-world applications. Additionally, understanding the mechanisms behind the antimicrobial actions of these extracts will be crucial for developing them into reliable therapeutic agents.

5. Conclusions

The study conclusively demonstrates the significant antimicrobial properties of garlic, black cumin seed oil, and honey, whether used individually or in combination, against various foodborne, opportunistic, and zoonotic infectious strains. The findings underscore the potential of these natural products as effective food additives and supplements, providing both nutritional and medicinal benefits. The study also highlights that solvent extraction, particularly with ethanol, is superior to non-solvent methods in enhancing antimicrobial activity. While the results are promising, further *in vivo* evaluations are necessary to bridge the gap between laboratory findings and real-world applications. The study's outcomes suggest that these natural remedies hold promise for developing new antimicrobial drugs, emphasizing the need for continued research, drug formulation testing, and policy development for the conservation and utilization of medicinal plants.

Acknowledgements

The author would like to acknowledge to the Jimma University College of Agriculture and Veterinary Medicine School of Veterinary Medicine, Jimma, Ethiopia.

Data availability

All relevant data are within the manuscript.

Informed consent statement

No informed consent was required to conduct the study.

Conflict of interest

The authors declare no conflict of interest.

Authors' contribution

Conceptualization: Hafiz Esmael; **Methodology:** Hafiz Esmael and Abdu Mohamed; **Formal analysis:** Hafiz Esmael; **Original draft preparation:** Hafiz Esmael and Kazi Abdus Sobur; **Manuscript formatting and result interpretation:** Md. Zaminur Rahman and

Md. Mosharrar Hossen; **First draft development, data analysis, review and editing:** Kazi Abdus Sobur and Md. Mosharrar Hossen; **Easing it, final review and editing the manuscript:** Md. Ashiquen Nobli.

References

- Abebe BA and Chane TS, 2021. Ethnobotanical study of medicinal plants used to treat human and livestock ailments in Hulet Eju Enese Woreda, east Gojjam zone of Amhara region, Ethiopia. *Evidence-Based Complementary and Alternative Medicine*, 2021: 6668541. <https://doi.org/0.1155/2021/6668541>
- Balkrishna A, Sharma N, Srivastava D, Kukreti A, Srivastava S and Arya V, 2024. Exploring the safety, efficacy, and bioactivity of herbal medicines: Bridging traditional wisdom and modern science in healthcare. *Future Integrative Medicine*, 3: 35-49. <https://doi.org/10.14218/FIM.2023.00086>
- Baynes RE, Dedonder K, Kissell L, Mzyk D, Marmulak T, Smith G and Riviere JE, 2016. Health concerns and management of select veterinary drug residues. *Food and Chemical Toxicology*, 88: 112-122. <https://doi.org/10.1016/j.fct.2015.12.020>
- Benkaci-Ali F, Baaliouamer A, Wathelet JP and Marlier M, 2012. Chemical composition and physicochemical characteristics of fixed oils from Algerian *Nigella sativa* seeds. *Chemistry of Natural Compounds*, 47: 925-931. <https://doi.org/10.1007/s10600-012-0106-7>
- Benkeblia N, 2004. Antimicrobial activity of essential oil extracts of various onions (*Allium cepa*) and garlic (*Allium sativum*). *LWT-food science and technology*, 37(2): 263-268. <https://doi.org/10.1016/j.lwt.2003.09.001>
- Chang Q, Wang W, Regev-Yochay G, Lipsitch M and Hanage WP, 2015. Antibiotics in agriculture and the risk to human health: how worried should we be?. *Evolutionary Applications*, 8(3): 240-247. <https://doi.org/10.1111/eva.12185>
- CLSI, 2020. Performance standards for antimicrobial susceptibility testing. Wayne State University Press, Detroit, MI, USA. pp. 1-332.
- Cushnie TPT and Lamb AJ, 2020. Antimicrobial activity of flavonoids. *International Journal of Antimicrobial Agents*, 26: 343-356. <https://doi.org/10.1016/j.ijantimicag.2005.09.002>
- Dessie AB, Abate TM, Adane BT, Tesfa T and Getu S, 2020. Estimation of technical efficiency of black cumin (*Nigella sativa* L.) farming in northwest Ethiopia: a stochastic frontier approach. *Journal of Economic Structures*, 9: 1-14. <https://doi.org/10.1186/s40008-020-00198-1>
- Durairaj S, Srinivasan S and Lakshmanaperumalsamy P, 2009. *In vitro* antibacterial activity and stability of garlic extract at different pH and temperature. *Electronic journal of Biology*, 5: 5-10.
- Gabriel T, Vestine A, Kim KD, Kwon SJ, Sivanesan I and Chun SC, 2022. Antibacterial activity of nanoparticles of garlic (*Allium sativum*) extract against different bacteria such as *Streptococcus mutans* and *Poryphomonas gingivalis*. *Applied Sciences*, 12(7): 3491. <https://doi.org/10.3390/app12073491>
- Hannan MA, Rahman MA, Sohag AAM, Uddin MJ, Dash R, Sikder MH and Kim B, 2021. Black cumin (*Nigella sativa* L.): A comprehensive review on phytochemistry, health benefits, molecular pharmacology, and safety. *Nutrients*, 13(6): 1784. <https://doi.org/10.3390/nu13061784>
- Hossain MS, Jony MAH, Saha N, Islam B, Sobur KA, Mowdood S and Hossain KM, 2024. Antibiotic resistance genes detection in *Escherichia coli* isolated from raw meat in Rajshahi division of Bangladesh. *American Journal of Microbiological Research*, 12(4): 79-84. <https://doi.org/10.12691/ajmr-12-4-1>
- Hutabarat RT, Nurjanah U and Fahrurrozi F, 2021. Effects of mulching on weed growth and cucumber yield. *Journal of Applied Horticulture*, 23: 125-129. <https://doi.org/10.37855/jah.2021.v23i02.24>

- Islam MA, Bose P, Rahman MZ, Muktaruzzaman M, Sultana P, Ahamed T and Khatun MM, 2024. A review of antimicrobial usage practice in livestock and poultry production and its consequences on human and animal health. *Journal of Advanced Veterinary and Animal Research*, 11(3): 675–685. <http://doi.org/10.5455/javar.2024.k817>
- Jadon R and Dixit S, 2014. Phytochemical extraction and antimicrobial activity of some medicinal plants on different microbial strains. *Journal of Medicinal Plants Studies*, 2: 58-63.
- Julqarnain SM, Bose P, Rahman MZ, Khatun MM and Islam MA, 2022. Bacteriological quality and prevalence of foodborne bacteria in broiler meat sold at live bird markets at Mymensingh City in Bangladesh. *Journal of Advanced Veterinary and Animal Research*, 9(3): 405. <http://doi.org/10.5455/javar.2022.i608>
- Magryś A, Olender A and Tchórzewska D, 2021. Antibacterial properties of *Allium sativum* L. against the most emerging multidrug-resistant bacteria and its synergy with antibiotics. *Archives of Microbiology*, 203(5): 2257-2268. <https://doi.org/10.1007/s00203-021-02248-z>
- Ncube B, Finnie JF and Van Staden J, 2012. In vitro antimicrobial synergism within plant extract combinations from three South African medicinal bulbs. *Journal of Ethnopharmacology*, 139: 81-89. <https://doi.org/10.1016/j.jep.2011.10.025>
- Okaiyeto K and Oguntibeju OO, 2021. African herbal medicines: Adverse effects and cytotoxic potentials with different therapeutic applications. *International Journal of Environmental Research and Public Health*, 18(11): 5988. <https://doi.org/10.3390/ijerph18115988>
- Petrosillo N, 2008. Natural products and wound management: a never-ending story. *Clinical Infectious Diseases: an Official Publication of the Infectious Diseases Society of America*, 47: 730-731. <https://doi.org/10.1086/590972>
- Radojević ID, Stanković MS, Stefanović OD, Topuzović MD and Čomić LR, 2012. Antioxidative and antimicrobial properties of different extracts from *Sideritis montana* L. *Romanian biotechnological letters*, 17(2): 7015.
- Rana AH, Bose P, Sobur KA, Hossen MM, Mowdood S, Hossain MK, Afroz F, Rumi NA, Hasan M, Jahan N, Titas AR and Hosen MA, 2024. Molecular detection and antibiogram profiles of *Escherichia coli* and *Vibrio cholerae* isolated from raw vegetables in the northern district of Bangladesh. *Journal of Bioscience and Environment Research*, 1: 26-34. <https://doi.org/10.69517/jber.2024.01.01.0006>
- Sadeghian A and Ghazvini K, 2015. Antimicrobial activity of garlic extract against *Shigella*. *Iranian Journal of Medical Sciences*, 27: 142-144.
- Saganuwan S, 2016. Competitive biodiversity of human and vascular plant species: the implications for pharmaceutical industries, health and world economy part-2. *European Journal of Medicinal Plants*, 16: 1-29. <https://doi.org/10.9734/EJMP/2016/27325>
- Sen T and Samanta SK, 2015. Medicinal plants, human health and biodiversity: a broad review. *Biotechnological Applications of Biodiversity*, 59-110. https://doi.org/10.1007/10_2014_273
- UNEP, 2015. Global Biodiversity assessment. United Nation Environmental Program. Addis Ababa, Ethiopia.
- Vaou N, Stavropoulou E, Voidarou C, Tsigalou C and Bezirtzoglou E, 2021. Towards advances in medicinal plant antimicrobial activity: A review study on challenges and future perspectives. *Microorganisms*, 9(10): 2041. <https://doi.org/10.3390/microorganisms9102041>
- WHO, 2014. Antimicrobial resistance: global report on surveillance. World Health Organization.
- Wolde T, Kuma H, Trueha K and Yabeker A, 2018. Anti-bacterial activity of garlic extract against human pathogenic bacteria. *Journal of Pharmacovigilance*, 6: 1-5. <https://doi.org/0.4172/2329-6887.1000253>
- Yimer EM, Tuem KB, Karim A, Ur-Rehman N and Anwar F, 2019. *Nigella sativa* L. (black cumin): a promising natural remedy for wide range of illnesses. *Evidence-Based Complementary and Alternative Medicine*, 2019: 1528635. <https://doi.org/10.1155/2019/1528635>
- Yineger H, Yewhalaw D and Teketay D, 2008. Plants of veterinary importance in Southwestern Ethiopia: The case of Gilgel Ghibe area. *Forests, Trees and Livelihoods*, 18: 165-181. <https://doi.org/10.1080/14728028.2008.9752627>
- Zhou X, Seto SW, Chang D, Kiat H, Razmovski-Naumovski V, Chan K and Bensoussan A, 2016. Synergistic effects of Chinese herbal medicine: a comprehensive review of methodology and current research. *Frontiers in Pharmacology*, 7: 201. <https://doi.org/10.3389/fphar.2016.00201>



Publisher's note

Genesis Publishing Consortium Limited pledges to maintain a neutral stance on jurisdictional claims shown in published maps and institutional affiliations.